

自己組織化粒子理論を用いた群れの研究

ラナ エムディー マスド 蔡 東生

鳥の群れ、陸上動物の群れ、魚の群れあるいは、歩行者の歩行は自然界の美しいものである。だが、動物のこの集合この集合的な行動のメカニズムに関して本わすかし知られていない。群れの中で個々の個体は群れ全体を意識せず、単純なローカルに従って行動する。本論文では、我々は群れの特徴を分析し、行動をシミュレーションする。本論文は主に鳥の群れにフォカスを与える。

Research of Aggregation Using Self-propelled Particles

RANA MD. MASUD DONGSHENG CAI

The aggregate motion of a flock of birds, a herd of land animals, a school of fish or even the walk of pedestrians are beautiful and familiar part of the natural world. But very few is known about the mechanism behind the emerging reason of aggregation. In aggregation individuals move interacting following simple local rules and do not consider the aggregation as a whole. In this paper, we will try to analysis the characteristics of individual in aggregation and simulate them. We will mainly focus on the collective behavior of birds in a flock.

1. Introduction

Collective behavior could be stated as “*the way in which an individual unit’s activity is dominated by its neighbors so that all units simultaneously alter their behavior to a common pattern*” [1]. By acting collectively, individuals (both organisms and non-living objects are considerable) synchronize their signals or motion. The main features of collective behavior are that an individual unit’s action is dominated by the influence of its neighbors – the unit behaves differently from the way it would behave on its own; and that such systems show interesting ordering phenomena as the units simultaneously change their behavior to a common pattern.

The aggregate motion of flock of birds, a herd of land animals, a school of fish are beautiful and nice examples of collective behavior. People clapping in phase during rhythmic applause, Mexican wave forming in stadia [2] also demonstrate collective behavior. Even non-living objects like ferromagnets show collective behavior. These materials can undergo spontaneous magnetization, in effect because they are made up of a host ‘tiny magnets’ [1].

Collective behavior of animals exhibits many contrasts. In case of flock of birds, flocks are made of discrete birds yet overall motion seems fluid; it is simple in concept yet is so visually complex, it seems randomly arrayed and yet is magnificently synchronized. The aggregation is constructed by the action of each individual, each action solely on the basis of its local perception of the world [3].

Scientists from different background involved themselves to understand and model different aggregations: school of fish [4], flock of birds [3, 5], pedestrian behavior [6]. Reynolds (1987) first introduced a bird flock model in computer graphics field [3]. He named the individual units as ‘boids’ related to ‘bird-like’ or ‘bird-oid’. To simulate a flock, he used three simple rules: (1) collision avoidance, (2) velocity matching and, (3) flock centering. Their simulation was confined to some tens to some hundreds individuals. These three rules seem reasonable, but they are unable to reproduce a flock once after the boids separate a little far away. Again, global consideration is not realistic.

Another simple model (SPP model) [7-9] showed that a individual need not to consider the whole flock to produce collective behavior. Only interactions with local neighbors and directional averaging with neighbors, while some environmental noise exists, is enough to produce collective motion. In their model, the individuals exist around a certain radius circle to a reference individual, are considered the neighbors of that reference individual. Therefore, collective behaviors created in this model greatly depend on density of the aggregation. However, recent field study from European scientists [10] confirmed that the starlings flock’s behavior is density independence. They argued that birds’ behavior depends on topological

distance rather than metric one.

In this paper, we tried to construct a bird flock of large numbers from some thousands to some ten thousands. We would base the basic *SPP model* for its simplicity [7], but include cohesion and collision avoidance. Though *SPP model* is strictly metric, we would exclude the metric perspective for individuals' interactions, instead, include the topological perspective while considering interactions with neighbors for the topological idea is supported from field study [10]. And we would try to extract and test some properties of flock of birds.

2. SPP Model

SPP stands for *Self-propelled particles*. The particles that make action or motion without the influence or action of any external force are called *self-propelled particle* [11]. In this sense, animals that produce collective behavior in different sort of aggregation, can be pointed as self-propelled particles. Instead of three rules model of Reynolds [3], SPP model [7] is based on only one rule: *at each time step, a given particle driven with a constant absolute velocity assumes the average direction of motion of the particles in its neighborhood of radius r with some random perturbation added*. The analogy can be formulated as follows: The rule corresponding to the ferromagnetic interaction tending to align the spins in the same direction is replaced by the rule of aligning the *direction of motion* of particles. Random perturbations are applied in analogy with the temperature. Biological subjects have the tendency to move as other subjects do in their neighborhood (Brien, 1989). Therefore, SPP model can be useful to model the flock of birds and other living organisms.

The simulations were carried out in a square shaped cell of linear size L with periodic boundary conditions. Interaction radius r was used as the unit to measure distances ($r = 1$), while the time unit, Δt was the time interval between two updating of direction and positions. The initial condition: (1) at time, $t = 0, N$ particles were randomly distributed in the cell, (2) had the same absolute velocity, v_0 and (3) randomly distributed direction. The velocities of particles $\{\vec{v}_i\}$ were determined at each time step, and the position of i th particle is updated according to-

$$\vec{v}_i(t + \Delta t) = v_0 \frac{\langle \vec{v}_j(t) \rangle_r}{|\langle \vec{v}_j(t) \rangle_r|} + \text{perturbation} \quad (1)$$

$$\vec{x}_i(t + \Delta t) = \vec{x}_i(t) + \vec{v}_i(t + \Delta t)\Delta t \quad (2)$$

Here $\langle \dots \rangle_r$ denotes averaging of the velocities within a circle of radius r surrounding

particle i . $\langle \vec{v}_j(t) \rangle_r / |\langle \vec{v}_j(t) \rangle_r|$ provides a unit vector pointing in the average direction of motion. Perturbation is taken account by adding a random angle corresponding to the average direction of motion in the neighborhood particle of i . Perturbations are random values taken from a uniform distribution in the interval of $[\eta\pi, \eta\pi]$. The only parameters of the model is the density -- the number of particles in unit square (for 2 dimensions) or unit volume (for 3 dimensions) -- the velocity, v_0 and the level of perturbation, $\eta < 1$. In two dimensional simulation, Vicsek showed that, for a wide velocity range ($0.003 < v_0 < 0.3$), and higher density ($\rho = 12.0$) and smaller level of noise or perturbation ($\eta = 0.1$), after some time steps, all particles move in the almost same direction i.e. synchronize themselves by locally interacting with each others.

In the *SPP Model*, Vicsek introduced a order parameter which denotes the level or ordered motion of the aggregation. The ordered parameter, ϕ , is determined as follows:

$$\phi = \frac{1}{Nv_0} \left| \sum_{i=1}^N \vec{v}_i \right| \quad (3)$$

Where N is the number of particles, \vec{v}_i is the velocity of i th particles. ϕ goes near to 1 when the aggregation is ordered and equal to 1 for fully ordered. In contrast, when ϕ is near to zero; it means that the particles are randomly walking and showing no collective behavior. .

3. Metric or Topology

Topological distance: The word „topology’ is derived from Greek word „topos’ which means place or space, and „logos’ which means study or idea or theory [13, 14]. Therefore topology can be understood as the study of place or space. “Topology,” the English form, was first used in 1883 in Listing's obituary in the journal Nature to distinguish "qualitative geometry from the ordinary geometry in which quantitative relations chiefly treated". In this paper, when we would talk about „metric distance’, we would mean the quantitative distance i.e. real distance. And when we use „topological distance’, we would rank the surrounding particles to a reference. The rank would be 1, for the most nearest neighbor, 2 for the second nearest neighbor and so on. These ranks would be the topological distances. Therefore topological distances would be discrete: 1, 2, 3,.. The important distinction is that topological distance does not change with the density of aggregation i.e. the most nearest neighbor’s rank would be 1 (topological distance = 1) no matter how far or how near it is. In economics, for example, the relevant quantity is not how many kilometers separate two countries (metric distance), but rather than the number of intermediate countries between them (topological distance) [15].

4. Ballerini's Field Study

Ballerini et al. (2008), by reconstructing three-dimensional positions of individual birds of few thousand members showed that the interactions among the birds do not depend on metric distance rather than depend on topological one. Moreover, each bird interacts with a fixed number of birds (6-7 birds). They tried to show that the topological interaction can achieve more cohesion than the metric one while robust cohesion is needed for complex density and shape changes of flock not breaking cohesion among birds..

The main goal of the interaction among individuals is to maintain cohesion of the aggregation. This is very strong biological requirement, shaped by the evolutionary pressure for survivor: stragglers and small groups are significantly more prone to predation than animals belonging to large and highly cohesive aggregation [16]. In topological model, cohesion among individuals does not vary with density changes, therefore more suitable to keep cohesion.

Ballerini et al. (2008) discussed about the characterization of structure of birds within flock is given by the spatial distribution of nearest neighbors. Given a reference bird, they measured the angular orientation of its nearest neighbors with respect to the flock's direction of motion. The measurement showed an anisotropic characteristic and the anisotropic characteristic tended to fade out as the rank of the nearest neighbors increases. This means that the anisotropic characteristics of flock is the result of individual interaction.

5. Results and Discussions

Ballerini et al. (2008) made a simple two dimensional predator-prey model based on *SPP model* to emphasize that the topological interaction should show strong cohesion. However, we reproduced the same results in two dimensional case and extended it to three dimensional predator-prey model. We have successfully shown that the three dimensional model exhibits the same type of cohesion as the two dimensional model does (Figure 1b and 1e).

(1) Predator-Prey Model

In predator-prey model (two dimensional), we used equation (1) and (2) to update prey's velocity and position. However, the perturbation or noise part is replaced by the impulsive force from the predator to prey. Predator's velocity and direction remain unchanged and does not have effect from preys. The impulsive force from predator to prey is determined as equation (4).

$$\vec{F}_i = f_0 \frac{\vec{r}_{predator-i}}{|\vec{r}_{predator-i}|^2} \quad (4)$$

\vec{F}_i is the impulsive force to i th bird, f_0 is the impulsive force posed by the predator and $\vec{r}_{predator-i}$ is the distance vector from predator to prey. For metric case, we used interaction radius as 0.15 and in case of topological situation, we assume that a bird interact with three nearest neighbor -- for two dimensional case individuals show optimum interaction when they interact with three nearest neighbors [4]. For both metric and topological case, we calculated the isolated individuals created by predator attack. Figure 1a shows that in metric case, maximum probability is for three isolated individuals while in topological case, Figure 1b shows that the maximum probability is for zero isolated individual. Again, the probability bars of separated individual decay very quickly in contrast with the metric interaction. Therefore, it shows that metric interaction is prone to topological interaction and topological interaction produces more cohesion among individuals in aggregation.

We can assume that the birds may have preference while aligning with the neighbors. We ran another simulation taking weighted average of neighbor's velocity. We modified equation (1) to equation (5) to update velocity, and found that cohesiveness increased (Figure 1d). We can not say for sure, but point out to birds may have preferences to among nearest neighbors. Same sort of characteristic has been achieved for three dimensional predator-prey simulation (Figure 1e).

$$\vec{v}_i(t + \Delta t) = v_0 \frac{\langle \vec{v}_j(t) \rangle_r}{|\langle \vec{v}_j(t) \rangle_r|} + perturbation \quad (5a)$$

$$\text{where, } \langle \vec{v}_j(t) \rangle_r = \frac{\vec{v}_i(t) + \sum_{j=1}^N \vec{v}_j(t) / (1+j)}{1 + \sum_{j=1}^N 1/(1+j)} \quad (5b)$$

(2) Density Independence

In topological interaction, interactions among individuals should be density independence, i.e. they should show the same sort of interaction results for different densities of aggregation. We have run simulations (the above two dimensional predator-prey model) for different densities and demonstrate that the characteristic of interaction vary negligibly. (Figure 2).

(3) Compatibility of SPP to Model

Is SPP is compatible to model bird flock? To test this, we have considered one of Ballerini's field study's result [10]. They defined a parameter called sparseness (r_1) – the

average first nearest neighbor distance of a flock – which is inverse proportion to the density of the flock; and metric range for topological interaction (r_c) – the average $n_c(= 7)$ th nearest neighbor distance of a flock – and found a strong linear correlation (Figure 3a) between them. We will take this as a test-stone to test the compatibility of SPP model. For ten different initial sparseness of our predator-prey model, we found that our simulations showed that there remains strong linear correlation between sparseness and metric range (Figure 3b).




Figure 1. The horizontal axis (in Figure (a, b, d and e)) shows the number isolated bird after attack and vertical axis shows the probability of that number isolated bird(s). In the model, we valued $v_0 = 0.25, f_0 = 0.05$. At $t = 0$, all birds are initialized with the same direction and the predator is at the opposite direction. Total individual is 200. Data is measured after 2000 time steps for simulation, and probability is taken after 2000 simulations done for each metric and topological case. The prey, initially are distributed a radius 1 circle and predator vertical position is 0.9 from the flock’s center. Interaction range for metric case, i.e. metric range is 0.15 and topological range is 3. We considered a bird is isolated if no other bird is present in 0.45 radius with respect to the reference bird. (a) shows the probability of isolated bird in metric case (maximum probability is 16.5% for 3 isolated bird), and (b) shows the probability in topological case, and the maximum probability is 52.4% for no isolated bird. (c) shows the image of the simulation; (d) Comparison between non-preferred and preferred velocity alignment. Preferred alignment shows better cohesion. (e) shows the simulation result for

three dimensional topology case. Time step is 1000, number of simulation is 1000. 1000 individuals, initially, are distributed in 1 unit radius sphere. The parameter values are, $v_0 = 0.25, f_0 = 0.05$, and isolation determination distance is 1.15.




Figure 2. Predator-prey model has been tested for different densities (different number of individuals are distributed within the same area). Other parameters coincide with the two dimensional topological model in section 5.1.




Figure 3. (a) Linear correlation between sparseness and metric range (Ballerini), Pearson correlation = 0.78. (b) Linear correlation between sparseness and metric range (simulation), Pearson correlation = 0.98.

(4) **Our Model**

In SPP model, we could produce some trend of flock’s behavior (staying together under perturbation and linear correlation between sparseness and metric range). But as only directional alignment has been considered, as time passes cohesion will break down [17]. In our predator-prey simulation, we found that even though boids have strong relation in alignment, the flock tends to get sparser as time passes even when there is negligible perturbation (Figure 4). Therefore, to model a flock consisting large number of individuals we have to consider some other interactive forces that are presented among individuals. Gruler et.

Al (1999) , and R. Kemkemer (2000) [18, 19] described that human melanocytes -- pigment cells of the skin -- are also act collectively without external force. That is why, melanocytes can be said as SPPs. But melanocytes do not show directional properties rather show apolar characteristics. Melanocytes show nematic arrangements (Figure 5) and their net motion is zero. They interact with each other nematically. This can be a vital interaction in different SPPs [11]. Vicsek model (1995) assumes objects as point like while melanocytes are as rod like. Therefore, to model bird flock, we can consider birds as a rod like objects that consider nematic forces for cohesion and also tend to make directional alignment. With this hypothesis, we will introduce a topological model where both nematic forces and tenderness for directional alignment would exist. By modifying SPP model with topological essence, we described the velocity update for each boid as equation (6).

$$\vec{v}_i(t + \Delta t) = v_0 v \{ s \sum_{j=1, i}^N \vec{v}_j(t) + (1 - s) \sum_{j=1}^N f_0 \vec{e}_{ij} + \eta \vec{z} \} \quad (6)$$




Figure 4. Sparseness increases with time steps.




Figure 5. Human melanocytes on a glass surface. We can see that these cells have nematic arrangements (Simha and Ramaswamt, 2002).

Here, v_0 is the speed, N is the number of neighbor for interaction, f_0 represents the nematic or cohesive force to each other, \vec{e}_{ij} is the unit vector to from i th boid to j th neighbor. η is the system's noise level, \vec{z} represents the random unit vector. \vec{v}_j is the velocity of j neighbor. s represents a strategy parameter, where, $0 \leq s \leq 1$. It determines to what extent, a bird is going to evaluate directional alignment and cohesion. Vicsek's (1995) SPP does not consider the prevention of collision among the individuals. We introduced collision prevention by imposing an infinite value to f_0 and, setting $\vec{e}_{ij} = -\vec{e}_{ji}$ when the nearest neighbor(s) are too close.

In large flocks, some characteristics can be found: density fluctuation, wave flow and complex patterns. SPP model for large number of particles shows density variance in the system both in two and three dimensions [17]. By simulating a large number of individuals with our proposed topological cohesive-directional alignment model, we were succeed to produce real like flock that showed density variation and complexity in patterns (Figure. 6). We argue that velocity alignment is responsible for density variation and nematic cohesive force is responsible for complex pattern [20]. However, yet, we have not been able to include wave flow in flock of birds. We are working on this.




Figure 6. A snapshot of flock of birds in our simulation. Number of individuals is 4096. Initially we distributed the individual randomly in a box of length 7 and initial directions were randomly taken. Individuals were updated according to equation (6) and equation (2). Time step was 1500. Other parameters are: $f_0 = 0.5$, $\eta = 0.001$, $s = 0.94$, $v_0 = 0.05$, $\Delta t = 1.0$, and collision prevention distance = 0.25.

6. Conclusion

We were able to produce a realistic structure of flock of birds using the topological interaction. We also could test and density independence characteristics of flock of birds that might be true. Still we are unable to create flat shape flocks and wave passing. We are working on this topic.

参考文献

- 1) Tamas Vicsek, "A question of scale," *Nature*, vol.411, (2001).
- 2) I. Farkas, D. Helbing and T. Vicsek, "Mexican waves in an excitable medium," *Nature*, 419, 131-132, (2002).
- 3) Craig W. Reynolds, "Flocks, Herds, and Schools: A Distributed Behavioral Model," *ACM Computer Graphics*, volume 21, No.4, (1987).
- 4) Yoshinobu Inada and Keiji Kawachi, "Order and Flexibility in the Motion of Fish Schools," *Journal of Theoretical Biology*, vol.214, issue 13, (2002).
- 5) K Bhattacharya and Tamas Vicsek, "Collective decision making in cohesive flocks," *New Journal of Physics*, 12 093019, (2010).
- 6) Mehdi Moussaïd, Dirk Helbing and Guy Theraulaz, "How simple rules determine pedestrian behavior and crowd disaster," *PNAS*, vol. 108, no. 17, (2011).
- 7) T. Vicsek, Andras Czirok, Eshel Ben-Jacob, Inon Cohen, and Ofer Shochet, "Novel Type of Phase Transition in a System of Self-Driven Particles," *Phys. Rev. Lett.* 75, 1226-1229, (1995).
- 8) B. Gönci, M. Nagy and T. Vicsek, "Phase transition in the scalar noise model of collective motion in three dimensions," *The European Physical Journal*, Volume 157, Number 1, 53-59, (2008).
- 9) Tamas Vicsek, "Universal Patterns of Collective Motion from Minimal Models of Flocking," 2008 Second IEEE International Conference on Self-Adaptive and Self-Organizing Systems, (2008).
- 10) M. Ballerini et al. "Interaction ruling animal collective behavior depends on topological rather than metric distance: Evidence from a field study," *PNAS*. vol. 105 no. 4 1232-1237, (2008).
- 11) R. Aditi Simha and Sriram Ramaswamy, "Statistical hydrodynamics of ordered suspensions of self-propelled particles: waves, giant number fluctuations and instabilities," *Physica A*. 306 262-269, (2002).
- 12) D. P. O'Brien, "Analysis of the internal arrangement of individuals within crustacean aggregations (Euphausiacea, Mysidacea)," *J. Exp. Mar. Biol. Ecol.* Vol. 128, pp. 1-30, (1989).
- 13) <http://en.wikipedia.org/wiki/Topology>. (26 September, 2011).
- 14) <http://www.nn.ij4u.or.jp/~hsat/techterm/topos.html> (26 September, 2011).
- 15) A. K. Henrikson, "Distance and foreign policy: a political geography approach," *Intl. Political Sci. Rev* 23, 437, (2002).
- 16) Ian Vine, "Risk of visual detection and pursuit by a predator and the selective advantage of flocking behaviour," *Journal of Theoretical Biology*, Volume 30, Issue 2, Pages 405-422, (1971).
- 17) H. Chaté, F. Ginelli, G. Grégoire, F. Peruani and F. Raynaud, "Modeling collective motion: variations on the Vicsek model," *The European Physical Journal B – Condensed Matter and Complex Systems*. Volume 64, Numbers 3-4, 451-456, (2008).
- 18) H. Gruler, U. Dewald and M. Eberhardt, "Nematic liquid crystals formed by living amoeboid cells," *The European Physical Journal B – Condensed Matter and Complex Systems*. Volume 11, Number 1, 187-192, (1999).
- 19) R. Kemkemer, D. Kling, D. Kaufmann and H. Gruler, "Elastic properties of nematoid arrangements formed by amoeboid cells," *The European Physical Journal E: Soft Matter and Biological Physics*. Volume 1, Numbers 2-3, 215-225, (2000).
- 20) F Heppner, "Three-dimensional structure and dynamics of bird flocks," *Animal Groups in Three Dimensions*, Julia K. Parrish, William Hamner, Cambridge University Press, (1997).