

Toric ideal の standard pair 分解と最小費用流問題

石関 隆幸[†] 今井 浩[‡]

[†]東京大学大学院理学系研究科情報科学専攻
113-0033 東京都文京区本郷 7-3-1
ishizeki@is.s.u-tokyo.ac.jp

[‡]東京大学大学院情報理工学系研究科コンピュータ科学専攻
113-0033 東京都文京区本郷 7-3-1
imai@is.s.u-tokyo.ac.jp

要旨

近年、整数計画問題に対してトーリックイデアルの離散性により、グレブナ基底や standard pair を用いた計算代数的アプローチが研究されている。これらのアプローチは既存の整数計画問題の解法に比べて計算量の改善を与えるものではないが、整数計画問題の構造について代数的な解析を与えている。本論文では、構造の知られている最小費用流問題に着目し、standard pair を用いた解析を行う。特に、トーリックイデアルのグレブナ基底および超幾何関数の結果を用いて、有向閉路を持たない有向グラフにおける最小費用流問題の(退化していない)双対多面体の頂点数の最小値が1になること、および最大値がカタラン数になることを示す。

Standard pair decompositions of toric ideals and minimum cost flow problems

Takayuki Ishizeki[†] and Hiroshi Imai[‡]

[†]Department of Information Science, Graduate School of Science, The University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
ishizeki@is.s.u-tokyo.ac.jp

[‡]Department of Computer Science, Graduate School of Information Science and Technology,
The University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan
imai@is.s.u-tokyo.ac.jp

Abstract

To integer programming problems, computational algebraic approaches using Gröbner bases or standard pairs via the discreteness of toric ideals have been studied in recent years. Although these approaches do not give improved time complexity bound compared with existing methods for solving integer programming problems, these give algebraic analysis of structure of integer programming problems. In this paper, we focus on the minimum cost flow problems, whose structure is well-known, and analyze using standard pairs. Especially, using results about Gröbner bases for toric ideals and hypergeometric functions, we show that the number of vertices of the (nondegenerate) dual polyhedra for minimum cost flow problems on acyclic directed graphs is more than 1 and less than the Catalan number.

1 Introduction

Recently, some algebraic approaches to integer programming problems have been studied. The two main approaches are using *Gröbner bases* [4] and *standard pair decompositions* [6]. Although they do not give improved complexity bounds compared with existing methods, these approaches themselves are very interesting by applying computational algebraic methods to such hard problems, and give algebraic analysis of structure of integer programming problems. The results in [4] and [6] have given several results about structure of integer programming problems using Gröbner bases [4, 11, 13, 14, 15] and standard pairs [6, 7].

The minimum cost network flow problem forms a well-known subclass of integer programming problems which can be solved in polynomial time. Gröbner basis approach for the minimum cost flow problems is a variant of cycle-canceling algorithm, in the sense that, for any feasible flow, we can obtain the optimal flow by augmenting flows along the cycles which correspond to the elements of Gröbner basis as many as possible. On the other hand, standard pair approach for the minimum cost flow problems first finds the set of standard pairs, and solves linear system of equations for each standard pair until an integer and non-negative solution is obtained. Since each standard pair for the minimum cost flow problem corresponds to a vertex of (nondegenerate) dual polyhedron of the problem, this approach is a variant of enumeration of dual bases.

The fact that each standard pair corresponds to a vertex of dual polyhedron of the minimum cost flow problem also implies that the number of standard pairs for the minimum cost flow problem gives the number of vertices of dual polyhedron of the problem. The number of vertices of dual polyhedra for some special problems have been studied. Especially for the transportation problem on $m \times n$ bipartite graph, which is equivalent to the minimum cost flow problem on the graph with $m + n$ vertices, the number of vertices is at most $\binom{m+n-2}{m-1}$ [2]. We show that, for the min-

imum cost flow problem on an *acyclic* directed graph with d vertices, the number of vertices of the dual polyhedron is at most $\frac{1}{2} \binom{2(d-1)}{d-1}$, the $(d-1)$ -th Catalan number. For the case of the problem on the acyclic tournament graph, there exists an example achieving this number. This upper bound is shown by combining two results, one is about the characterization of Gröbner basis for minimum cost flow problem [8], and the other is about the special hypergeometric function [5].

This paper is organized as follows. In Section 2, standard pair decompositions are defined and their relation with regular triangulations, dual polyhedra and integer programming problems are introduced. In Section 3, we characterize the standard pairs for the minimum cost flow problems. In Section 4, the minimum and the maximum number of vertices of dual polyhedra are shown for the minimum cost flow problems on acyclic tournament graphs. The methods which are used to show these results are also shown. For the problems on acyclic directed graphs, these results give upper and lower bounds. Finally in Section 5 we conclude this paper.

2 Standard pair decompositions of integer programs

In this section, we give some definitions about standard pairs, and their relation to regular triangulations, dual polyhedra and integer programmings. We refer to [12] for the introduction of standard pairs, and [6, 7] for their applications.

2.1 Standard pairs, regular triangulations and dual polyhedra

We fix a matrix $A \in \mathbb{Z}^{d \times n}$ and a cost vector $c \in \mathbb{R}^n$, and consider the family $IP_{A,c}$ of integer programming problems $IP_{A,c}(b) := \text{minimize}\{c \cdot x \mid Ax = b, x \in \mathbb{N}^n\}$ as b varies in $\{Au \mid u \in \mathbb{N}^n\} \subseteq \mathbb{Z}^d$ (\mathbb{N} is the set of non-negative integers, and \mathbb{Z} is the set of integers). We assume that c is *generic* in the sense that each program in $IP_{A,c}$ has a unique optimal solution. For a fixed c , let $O_c \subset \mathbb{N}^n$ be the

set of all the optimal solutions to all programs in $IP_{A,c}$, and $N_c := \mathbb{N}^n \setminus O_c$. Then there exists a unique minimal finite set $\{p_1, \dots, p_s\} \subset \mathbb{N}^n$ such that $N_c = \bigcup_{i=1}^s (p_i + \mathbb{N}^n)$, where $p_i + \mathbb{N}^n := \{p_i + u \mid u \in \mathbb{N}^n\}$ [13]. Algebraically, p_1, \dots, p_s are the set of exponent vectors of initial terms of G_c , the reduced Gröbner bases of the toric ideal I_A with respect to c .

For $a \in \mathbb{N}^n$ and $\sigma \subseteq \{1, \dots, n\}$, we define a set of points (a, σ) in \mathbb{N}^n as $(a, \sigma) := \{a + \sum_{i \in \sigma} k_i e_i \mid k_i \in \mathbb{N}\}$, where e_i is a i -th unit vector in \mathbb{R}^n .

Definition 2.1 (a, σ) is a standard pair of O_c if

- (i) the support of a is contained in $\{1, \dots, n\} \setminus \sigma$,
- (ii) $(a, \sigma) \subseteq O_c$, and
- (iii) $(a, \sigma) \not\subseteq (b, \tau)$ for any other (b, τ) which satisfies (i) and (ii).

We denote $S(O_c)$ the set of all standard pairs of O_c .

The standard pairs of O_c induce a unique covering of O_c , which we call the *standard pair decomposition* of O_c . $|S(O_c)|$ is called the *arithmetic degree* of O_c and denoted by $\text{arith-deg}(O_c)$. Because of the genericity of c , O_c is not empty and $\text{arith-deg}(O_c) \geq 1$.

Example 2.2 Let A be the incidence matrix of the network shown in Figure 1. For $c = (3, 1, 2)$, the standard pairs of $O_{(3,1,2)}$ are $((0, 0, 0), \{1, 2\})$ and $((0, 0, 0), \{2, 3\})$, thus the arithmetic degree of $O_{(3,1,2)}$ is 2. On the other hand, for $c = (1, 4, 2)$, the standard pair of $O_{(1,4,2)}$ is $((0, 0, 0), \{1, 3\})$, thus the arithmetic degree of $O_{(1,4,2)}$ is 1.

Let $\{a_1, \dots, a_n\}$ be the column vectors of A and $\text{cone}(A)$ the cone generated by a_1, \dots, a_n . For $\sigma \subset \{1, \dots, n\}$, we denote A_σ for the submatrix of A whose columns are indexed by σ . For a cost vector c , we define the *regular triangulation* Δ_c of $\text{cone}(A)$ as follows: $\text{cone}(A_\sigma)$ is a face of Δ_c if and only if there exists a vector $y \in \mathbb{R}^d$ such that




Figure 1: The network N and two types of standard pair decompositions

$y \cdot a_j = c_j$ ($j \in \sigma$) and $y \cdot a_j < c_j$ ($j \notin \sigma$). If $\text{cone}(A_\sigma)$ is a face of Δ_c , σ also is called a *face* of Δ_c . The genericity of c implies that Δ_c is in fact a triangulation (i.e. each face of Δ_c is simplicial) [11].

Lemma 2.3 ([10, 12]) (i) If O_c has $(*, \sigma)$ as a standard pair, then σ is a face of Δ_c .

(ii) O_c has $((0, \dots, 0), \sigma)$ as a standard pair if and only if σ is a maximal face of Δ_c .

(iii) If a_1, \dots, a_n span an affine hyperplane, then Δ_c is same as a regular triangulation of $\text{conv}(A)$ with respect to c , and the number of standard pairs $(*, \sigma)$ for a maximal face σ of Δ_c equals the normalized volume of σ in Δ_c .

When the vertices of $\text{conv}(A)$ are in the m -dimensional lattice $L \simeq \mathbb{Z}^m$, we define the *normalized volume* of a maximal face σ of Δ_c by the volume of σ with the normalization that the volume of the convex hull of $0, e_1, \dots, e_m$ is 1. Here, $\{e_i\}_{1 \leq i \leq m}$ are the basis of the lattice L .

For a polyhedron $P \subset \mathbb{R}^n$ and a face F of P , the *normal cone* of F at P is the cone $N_P(F) := \{\omega \in \mathbb{R}^n \mid \omega \cdot x' \geq \omega \cdot x \text{ for all } x' \in F \text{ and } x \in P\}$. The set of normal cones for all faces of P is called the *normal fan* of P .




Figure 2: The dual polyhedron $P_{(3,1,2)}$ and the triangulation $\Delta_{(3,1,2)}$ of Example 2.2

Lemma 2.4 ([7]) Δ_c is the normal fan of the polyhedron $P_c := \{y \in \mathbb{R}^d \mid yA \leq c\}$.

We remark that P_c is the dual polyhedron of the linear relaxation problem

$$LP_{A,c}(b) := \text{minimize } \{c \cdot x \mid Ax = b, x \geq 0\}$$

of $IP_{A,c}(b)$. This lemma shows that there is one-to-one correspondence between the vertices of P_c and the maximal faces of Δ_c .

Example 2.2 (continued.) Figure 2 shows the dual polyhedron $P_{(3,1,2)}$ and $\Delta_{(3,1,2)}$. We remark that $\text{cone}(A)$ is contained in the hyperplane $x_1 + x_2 + x_3 = 0$.

2.2 Standard pair decompositions and integer programs

Using the standard pair decomposition of O_c , $IP_{A,c}(b)$ can be solved by solving at most (arithmetic degree of O_c)-many linear systems of equations [6]. We describe the outline of this approach.

Let u be the optimal solution to $IP_{A,c}(b)$. Since the standard pairs cover O_c , u is covered by some standard pair (a, σ) . Thus $u = a + \sum_{i \in \sigma} k_i e_i$ for some non-negative integers $\{k_i\}_{i \in \sigma}$, and

$$\begin{aligned} b &= Au \\ &= A \left(a + \sum_{i \in \sigma} k_i e_i \right) \\ &= Aa + \sum_{i \in \sigma} k_i a_i. \end{aligned}$$

Lemma 2.3 implies that $\{a_i\}_{i \in \sigma}$ are linearly independent, and therefore $\{k_i\}_{i \in \sigma}$ is the unique solution to the linear system

$$\sum_{i \in \sigma} x_i a_i = b - Aa. \quad (1)$$

This observation induces an algorithm to solve $IP_{A,c}(b)$ using the standard pair decomposition of O_c .

Algorithm 2.3 (Solving $IP_{A,c}(b)$ using $S(O_c)$)

- (i) For $(a, \sigma) \in S(O_c)$, solve the linear system (1). Let $\{k_i\}_{i \in \sigma}$ be the solution.
- (ii) If $\{k_i\}_{i \in \sigma}$ are all both integral and non-negative, output $a + \sum_{i \in \sigma} k_i e_i$ as the optimal solution. Otherwise, repeat (i) for another standard pair.

This algorithm solves at most $\text{arith-deg}(O_c)$ -many linear systems of equations. Therefore arithmetic degree of O_c is a measure of the complexity of $IP_{A,c}$.

3 Standard pairs of directed graphs

We consider that A is the incidence matrix of a network $N = (V, E)$ such that $|V| = d$, $|E| = n$, $b \in \{Au \mid u \in \mathbb{N}^n\}$ and $c \in \mathbb{R}_{\geq 0}^n$. This means that $IP_{A,c}(b)$ is an uncapacitated minimum cost flow problem on N with the demand/supply vector b . In this case, N_c has a special property as follows.

Lemma 3.4 Let $\{p_1, \dots, p_s\}$ be the minimal vectors such that $N_c = \bigcup_{i=1}^s (p_i + \mathbb{N}^n)$. Then all p_1, \dots, p_s are 0-1 vectors for any cost vector $c \in \mathbb{R}_{\geq 0}^n$.

Proof: In [8], we showed that reduced Gröbner basis for I_A with respect to any c corresponds to the set of circuits in N . This implies that the exponent vectors of reduced Gröbner basis are 0-1 vectors. \square

When N_c is generated by 0-1 vectors, the set $S(O_c)$ are obtained by all of the maximal faces of Δ_c .

Lemma 3.5 ([7]) Let $\{p_1, \dots, p_s\}$ be the minimal vectors such that $N_{\mathbf{c}} = \bigcup_{i=1}^s (p_i + \mathbb{N}^n)$. Then p_1, \dots, p_s are all 0-1 vectors if and only if $S(O_{\mathbf{c}}) = \{(0, \dots, 0), \sigma\} \mid \sigma \text{ is the maximal faces of } \Delta_{\mathbf{c}}\}$.

The edges in the optimum flow of uncapacitated minimum cost flow problem define a forest [1]. Therefore, with the fact the dimension of $\text{cone}(A)$ equals $d - 1$, the next proposition is implied by Lemma 2.3, 3.4 and 3.5.

Proposition 3.6 Any standard pair of $O_{\mathbf{c}}$ is $((0, \dots, 0), \sigma)$, where $\sigma \subseteq \{1, \dots, n\}$ and σ is a spanning tree of the network N .

Since there is one-to-one correspondence between the standard pairs $((0, \dots, 0), *)$ of $O_{\mathbf{c}}$ and the vertices of dual polyhedron, Algorithm 2.3 for the minimum cost flow problem $IP_{A, \mathbf{c}}(\mathbf{b})$ is a variant of the enumeration of dual feasible bases.

4 Arithmetic degree of acyclic directed graphs

In the case that the network is the acyclic tournament graph, the minimum and maximum arithmetic degrees can be obtained using some results about Gröbner bases. When the network is an acyclic directed graph, these results give upper and lower bounds for the arithmetic degree. Since each standard pair of the minimum cost flow problem corresponds to the vertex of the dual polyhedron for the linear relaxation of the problem, these results also give upper and lower bounds for the number of vertices of the dual polyhedron. In this section we show that, for the acyclic tournament graph with d vertices, the minimum arithmetic degree is 1 and the maximum arithmetic degree is $\frac{1}{d} \binom{2(d-1)}{d-1}$, which is the $(d-1)$ -Catalan number.

We assume that the vertices of acyclic directed graph have labels $1, \dots, d$ such that each edge (i, j) is oriented from the vertex i to the vertex j . We denote c_{ij} the cost of the edge (i, j) and x_{ij} the variable corresponds to

the edge (i, j) . The number of edges equals $n = \binom{d}{2}$.

4.1 Minimum arithmetic degree of acyclic tournament graphs

We remark again that the genericity of \mathbf{c} implies that, the arithmetic degree of $O_{\mathbf{c}}$ is equal to or greater than 1. We show that there exists a cost vector \mathbf{c} for which the arithmetic degree equals 1 for any d .

Theorem 4.1 For the acyclic tournament graph with d vertices, the minimum arithmetic degree of $O_{\mathbf{c}}$ which \mathbf{c} varies all generic cost vectors equals 1.

To show this theorem, we use a characterization of one $N_{\mathbf{c}}$.

Lemma 4.2 When $\mathbf{c} = (c_{ij})_{1 \leq i < j \leq d}$ satisfies that $c_{ij} > c_{i,i+1} + c_{i+1,i+2} + \dots + c_{j-1,j}$ for any $i < j - 1$, then $N_{\mathbf{c}} = \bigcup_{i=1}^{d-1} (e_{i,i+1} + \mathbb{N}^n) \subset \mathbb{N}^n$.

Proof: In [8], we showed that reduced Gröbner basis $G_{\mathbf{c}}$ for I_A with respect to the above \mathbf{c} is $\{x_{ij} - x_{i,i+1}x_{i+1,i+2} \dots x_{j-1,j} \mid 1 \leq i < j - 1 \leq d - 1\}$. Especially, the set of initial terms of $G_{\mathbf{c}}$ equals $\{x_{ij} \mid 1 \leq i < j - 1 \leq d - 1\}$, which implies that $N_{\mathbf{c}} = \bigcup_{i=1}^{d-1} (e_{i,i+1} + \mathbb{N}^n)$. \square

Proof of Theorem 4.1: Because of Lemma 4.2, $\mathbf{a} = (a_{ij})_{1 \leq i < j \leq d} \in \mathbb{N}^n$ is in $O_{\mathbf{c}}$ for \mathbf{c} as in Lemma 4.2 if and only if $a_{ij} = 0$ for any (i, j) such that $j - i > 1$. The set of all such points coincides $((0, \dots, 0), \{(1, 2), (2, 3), \dots, (d - 1, d)\})$. Thus only this pair is a standard pair of $O_{\mathbf{c}}$. \square

4.2 Maximum arithmetic degree of acyclic tournament graphs

To show the maximum arithmetic degree, we use another integer programming problem defined from the original integer programming problem. For a general matrix $A \in \mathbb{Z}^{d \times n}$, we

define another matrix $A' \in \mathbb{Z}^{(d+1) \times (n+1)}$ as

$$A' := \begin{pmatrix} 1 & 1 & \cdots & 1 & 1 \\ & & & & 0 \\ & & A & & \vdots \\ & & & & 0 \end{pmatrix} = \begin{pmatrix} 1 & 1 & \cdots & 1 & 1 \\ & & & & 0 \\ \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n & \vdots \\ & & & & 0 \end{pmatrix}. \quad (2)$$

Let $\mathbf{a}'_i = \begin{pmatrix} 1 \\ \mathbf{a}_i \end{pmatrix}$ for $1 \leq i \leq n$ and \mathbf{a}'_{n+1} be the $n+1$ -th column vector of A' . We remark that $\mathbf{a}'_1, \dots, \mathbf{a}'_n, \mathbf{a}'_{n+1}$ span an affine hyperplane.

We define another family $IP_{A',(\mathbf{c},0)}$ of integer programming problem

$$IP_{A',(\mathbf{c},0)}(\mathbf{b}, \beta) := \text{minimize } \left\{ \mathbf{c} \cdot \mathbf{x} \mid \begin{array}{l} A'(\mathbf{x}_{n+1}) = \begin{pmatrix} \beta \\ \mathbf{b} \end{pmatrix}, \\ (\mathbf{x}_{n+1}) \in \mathbb{N}^{n+1} \end{array} \right\}$$

as $\begin{pmatrix} \beta \\ \mathbf{b} \end{pmatrix}$ varies in $\{A'\mathbf{u} \mid \mathbf{u} \in \mathbb{N}^{n+1}\}$. Similarly as $O_{\mathbf{c}}$, let $O'_{(\mathbf{c},0)} \subset \mathbb{N}^{n+1}$ be the set of all optimal solutions to all programs in $IP_{A',(\mathbf{c},0)}$, and $N'_{(\mathbf{c},0)} := \mathbb{N}^{n+1} \setminus O'_{(\mathbf{c},0)}$.

Proposition 4.3 (\mathbf{a}, σ) ($\mathbf{a} \in \mathbb{N}^n$, $\sigma \subseteq \{1, \dots, n\}$) is a standard pair of $O_{\mathbf{c}}$ if and only if $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right)$ is a standard pair of $O'_{(\mathbf{c},0)}$.

Proof: We first show that $(\mathbf{a}, \sigma) \subseteq O_{\mathbf{c}}$ if and only if $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right) \subseteq O'_{(\mathbf{c},0)}$. Suppose that $(\mathbf{a}, \sigma) \subseteq O_{\mathbf{c}}$ and choose any $\begin{pmatrix} \mathbf{u} \\ k \end{pmatrix} \in \left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right)$. If there exist any other $v \in \mathbb{N}^n$ and non-negative integer l such that $A'(\begin{pmatrix} v \\ l \end{pmatrix}) = A'(\begin{pmatrix} \mathbf{u} \\ k \end{pmatrix})$ and $\begin{pmatrix} v \\ l \end{pmatrix} \neq \begin{pmatrix} \mathbf{u} \\ k \end{pmatrix}$, then $A\mathbf{u} = A\mathbf{v}$, and $\mathbf{c} \cdot \mathbf{u} < \mathbf{c} \cdot \mathbf{v}$ since $\mathbf{u} \in O_{\mathbf{c}}$. Therefore, $\begin{pmatrix} \mathbf{u} \\ k \end{pmatrix}$ is the optimal solution to $IP_{A',(\mathbf{c},0)}(A\mathbf{u}, \beta)$ with $\beta = \sum_{i=1}^n u_i$. If there does not exist such $\begin{pmatrix} v \\ l \end{pmatrix}$, then clearly $\begin{pmatrix} \mathbf{u} \\ k \end{pmatrix}$ is the optimal for this integer programming problem. This shows that $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right) \subseteq O'_{(\mathbf{c},0)}$.

Conversely, suppose that $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right) \subseteq O'_{(\mathbf{c},0)}$ and choose any $\mathbf{u} \in (\mathbf{a}, \sigma)$. If there exists some $v \in \mathbb{N}^n$

such that $A\mathbf{v} = A\mathbf{u}$, then $A'(\begin{pmatrix} v \\ p \end{pmatrix}) = A'(\begin{pmatrix} \mathbf{u} \\ q \end{pmatrix})$ for any non-negative integers p, q such that $p - q = \sum_{i=1}^n v_i - \sum_{i=1}^n u_i$. Since $\begin{pmatrix} \mathbf{u} \\ p \end{pmatrix} \in \left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right) \subseteq O'_{(\mathbf{c},0)}$, and $(\mathbf{c}, 0) \cdot \begin{pmatrix} \mathbf{u} \\ p \end{pmatrix} < (\mathbf{c}, 0) \cdot \begin{pmatrix} v \\ q \end{pmatrix}$, which imply that $\mathbf{c} \cdot \mathbf{u} < \mathbf{c} \cdot \mathbf{v}$. Therefore, \mathbf{u} is the optimal solution to $IP_{A,\mathbf{c}}(A\mathbf{u})$. If there does not exist such v , then clearly \mathbf{u} is the optimal for this integer programming problem. Thus $(\mathbf{a}, \sigma) \subseteq O_{\mathbf{c}}$.

If $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right)$ is a standard pair of $O'_{(\mathbf{c},0)}$, then $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right) \not\subseteq \left(\begin{pmatrix} \mathbf{a}' \\ 0 \end{pmatrix}, \tau \cup \{n+1\}\right)$ for any other $\left(\begin{pmatrix} \mathbf{a}' \\ 0 \end{pmatrix}, \tau \cup \{n+1\}\right)$ which satisfies the condition (i) and (ii) in Definition 2.1 for $O'_{(\mathbf{c},0)}$. Since if $(\mathbf{a}, \sigma) \subset (\mathbf{a}', \tau)$ for any other (\mathbf{a}', τ) which satisfies (i) and (ii) in Definition 2.1 for $O_{\mathbf{c}}$, then $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right)$ must be contained in $\left(\begin{pmatrix} \mathbf{a}' \\ 0 \end{pmatrix}, \tau \cup \{n+1\}\right)$, which contradicts the assumption. Thus (\mathbf{a}, σ) is a standard pair of $O_{\mathbf{c}}$.

On the other hand, if $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right)$ is not a standard pair of $O'_{(\mathbf{c},0)}$, then there exists some $\left(\begin{pmatrix} \mathbf{a}' \\ k \end{pmatrix}, \tau'\right)$ such that $\left(\begin{pmatrix} \mathbf{a}' \\ k \end{pmatrix}, \tau'\right)$ contains $\left(\begin{pmatrix} \mathbf{a} \\ 0 \end{pmatrix}, \sigma \cup \{n+1\}\right)$ and $\left(\begin{pmatrix} \mathbf{a}' \\ k \end{pmatrix}, \tau'\right)$ satisfies (i) and (ii) in Definition 2.1 for $O'_{(\mathbf{c},0)}$. Then $n+1 \in \tau'$, and therefore $k = 0$ by (i) in Definition 2.1. Therefore, (\mathbf{a}', τ) where $\tau := \tau' \setminus \{n+1\}$ contains (\mathbf{a}, σ) and satisfies (i) and (ii) in Definition 2.1. Thus (\mathbf{a}, σ) is not a standard pair of $O_{\mathbf{c}}$. This completes the proof. \square

Example 2.2 (continued.) For this A , enlarged matrix A' is

$$A' = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \\ 0 & -1 & -1 & 0 \end{pmatrix}.$$

For $\mathbf{c} = (3, 1, 2)$, the standard pairs of $O'_{(3,1,2,0)}$ are $\left(\begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \{1, 2, 4\}\right)$ and $\left(\begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \{2, 3, 4\}\right)$. On the other hand, for $\mathbf{c} = (1, 4, 2)$, the standard pairs of $O'_{(1,4,2,0)}$ are $\left(\begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \{1, 2, 3\}\right)$ and

$((0, 0, 0, 0), \{1, 3, 4\})$. In this case, the only standard pair $((0, 0, 0, 0), \{1, 3, 4\})$ satisfies the condition in Proposition 4.3, which corresponds to the standard pair $((0, 0, 0), \{1, 3\})$ of $O_{(1,4,2)}$.

Theorem 4.4 For the acyclic tournament graph with d vertices, the maximum arithmetic degree of $O_{\mathbf{c}}$ which \mathbf{c} varies all generic cost vectors equals

$$C_{d-1} := \frac{1}{d} \binom{2(d-1)}{d-1},$$

which is the $(d-1)$ -th Catalan number.

To show this theorem, we prepare some lemmas. The first is by Gelfand et al. [5] which studies about some hypergeometric function.

Lemma 4.5 ([5]) Let A' be the enlarged matrix (2) for the incidence matrix A of the acyclic tournament graph with d vertices, and $\text{conv}(A')$ be the convex hull of $\mathbf{a}'_1, \dots, \mathbf{a}'_{n+1}$. Then the normalized volume of $\text{conv}(A')$ equals the $(d-1)$ -th Catalan number C_{d-1} .

Given a regular triangulation $\Delta'_{(\mathbf{c},k)}$ of $\text{conv}(A')$, the normalized volume of $\text{conv}(A')$ equals the sum of normalized volume for all maximal faces in $\Delta'_{(\mathbf{c},k)}$. Since the column vectors in $\mathbf{a}'_1, \dots, \mathbf{a}'_{n+1}$ span an affine hyperplane, the normalized volume of $\text{conv}(A')$ gives the number of standard pairs of $O'_{(\mathbf{c},k)}$ which correspond to the maximal faces of $\Delta'_{(\mathbf{c},k)}$ by Lemma 2.3 (iii).

Lemma 4.6 When $\mathbf{c} = (c_{ij})_{1 \leq i < j \leq d}$ satisfies that

- $c_{ij} + c_{jk} > c_{ik}$ for any $i < j < k$ and
- $c_{ik} + c_{jl} > c_{il} + c_{jk}$ for any $i < j < k < l$,

then

$$\begin{aligned} N_{\mathbf{c}} &= \bigcup_{1 \leq i < j < k \leq d} ((e_{ij} + e_{jk}) + \mathbb{N}^n) \\ &\cup \bigcup_{1 \leq i < j < k < l \leq d} ((e_{ik} + e_{jl}) + \mathbb{N}^n) \subset \mathbb{N}^n. \end{aligned}$$

Proof: In [8], we showed that reduced Gröbner basis $G_{\mathbf{c}}$ for I_A with respect to the above \mathbf{c} is $\{x_{ij}x_{jk} - x_{ik} \mid 1 \leq i < j < k \leq d\} \cup \{x_{ik}x_{jl} - x_{il}x_{jk} \mid 1 \leq i < j < k < l \leq d\}$. Especially, the set of initial terms of $G_{\mathbf{c}}$ equals $\{x_{ij}x_{jk} \mid 1 \leq i < j < k \leq d\} \cup \{x_{ik}x_{jl} \mid 1 \leq i < j < k < l \leq d\}$, which implies that $N_{\mathbf{c}} = \bigcup_{1 \leq i < j < k \leq d} ((e_{ij} + e_{jk}) + \mathbb{N}^n) \cup \bigcup_{1 \leq i < j < k < l \leq d} ((e_{ik} + e_{jl}) + \mathbb{N}^n)$. \square

Proof of Theorem 4.4: Because of Proposition 3.6, any standard pair of $O_{\mathbf{c}}$ is $((0, \dots, 0), \sigma)$ for any \mathbf{c} , and corresponds to the standard pair $((0, \dots, 0, 0), \sigma \cup \{n+1\})$ of $O'_{(\mathbf{c},0)}$. Especially, $\sigma \cup \{n+1\}$ is a maximal face of $\Delta'_{(\mathbf{c},0)}$. Therefore,

$$\begin{aligned} \text{arith-deg}(O_{\mathbf{c}}) &= |\{((0, \dots, 0), \sigma) \in S(O_{\mathbf{c}})\}| \\ &= \left| \{((0, \dots, 0), \sigma \cup \{n+1\}) \in S(O'_{(\mathbf{c},0)})\} \right| \\ &\leq \sum_{\tau} \left| \{(*, \tau) \in S(O'_{(\mathbf{c},0)})\} \right| \quad (3) \\ &= \text{normalized volume of } \text{conv}(A') \\ &= C_{d-1} \end{aligned}$$

where the sum in (3) is taken over all maximal faces τ of $\Delta'_{(\mathbf{c},0)}$.

Because of Proposition 3.6 and Lemma 4.6, for \mathbf{c} as in Lemma 4.6, $((0, \dots, 0), \sigma)$ is a standard pair of $O_{\mathbf{c}}$ if and only if σ is a spanning tree of the acyclic tournament graph which satisfies the following two conditions:

- (A) there are no $1 \leq i < j < k \leq d$ such that both (i, j) and (j, k) are edges in σ , and
- (B) there are no $1 \leq i < j < k < l \leq d$ such that both (i, k) and (j, l) are edges in σ .

The number of spanning trees of the complete graph with d vertices which satisfies (A) and (B) are known to be equal to the $(d-1)$ -th Catalan number (e.g. see [9]). \square

We remark that the Catalan number equals $\frac{4^n}{\sqrt{\pi n^{3/2}}} (1 + O(\frac{1}{n}))$ (e.g. see [3]).

Example 2.2 (continued.) $\mathbf{c} = (3, 1, 2)$ gives an example achieving the maximum arithmetic degree since $C_2 = \frac{1}{3} \binom{4}{2} = 2$. On the other hand, $\mathbf{c} = (1, 4, 2)$ gives an example achieving the minimum arithmetic degree.

4.3 Arithmetic degree of acyclic directed graphs

Since any acyclic directed graph G with d vertices can be seen as the subgraph of acyclic tournament graph \tilde{G} with same number of vertices by deleting some edges, the incidence matrix A of G is a submatrix of that \tilde{A} of \tilde{G} by deleting the columns which correspond to the deleted edges. Let A' (resp. \tilde{A}') be the enlarged matrix (2) of A (resp. \tilde{A}). Then the normalized volume of $\text{conv}(A')$ is equal to or less than that of $\text{conv}(\tilde{A}')$, which implies that the arithmetic degree for $IP_{A,c}$ for any cost vector c is equal to or less than the maximum arithmetic degree for the acyclic tournament graph, which is the $(d-1)$ -th Catalan number.

5 Conclusion

In this paper, we showed the bounds for the number of vertices of the dual polyhedra for the minimum cost flow problems on the acyclic directed graphs, using the results from Gröbner bases and hypergeometric functions. We also showed two examples for the acyclic tournament graphs, one achieves the minimum arithmetic degree 1, and the other achieves the maximum arithmetic degree C_{d-1} .

On the other hand, the arithmetic degrees for cyclic directed graphs are not known. Similar approach may be able to attack to this analysis, which should be a future work.

References

- [1] R. K. Ahuja, T. L. Magnanti and J. B. Orlin. *Network Flows: Theory, Algorithms, and Applications*. Prentice Hall, New Jersey, 1993.
- [2] M. L. Balinski and A. Russakoff. Faces of dual transportation polyhedra. *Mathematical Programming Study*, **22**(1984), pp. 1-8.
- [3] T. H. Cormen, C. E. Leiserson and R. L. Rivest. *Introduction to Algorithms*. MIT Press, Cambridge, MA, 1990.
- [4] P. Conti and C. Traverso. Buchberger Algorithm and Integer Programming. In *Proceedings of the ninth Applied Algebra, Algebraic Algorithms and Error-Correcting Codes (AAECC-9)* (New Orleans), Springer, LNCS **539**(1991), pp. 130-139.
- [5] I. M. Gelfand, M. I. Graev and A. Postnikov. Combinatorics of hypergeometric functions associated with positive roots. In *Arnold-Gelfand Mathematical Seminars: Geometry and Singularity Theory*, Birkhäuser, Boston, 1996, pp. 205-221.
- [6] S. Hoçten and R. R. Thomas. Standard pairs and group relaxations in integer programming. *Journal of Pure and Applied Algebra*, **139**(1999), pp. 133-157.
- [7] S. Hoçten and R. R. Thomas. Gomory integer programs. Los Alamos e-print archive, math.OC/0106031, 2001.
- [8] T. Ishizeki and H. Imai. Gröbner Bases of Acyclic Directed Graphs and Minimum Cost Flow Problems. *Proceedings of the 2nd Japanese-Hungarian Symposium on Discrete Mathematics and Its Applications*, Budapest, pp. 82-91, 2001.
- [9] R. Stanley. *Enumerative Combinatorics Vol. 2*. Cambridge Studies in Advanced Mathematics, Vol. 62, Cambridge University Press, Cambridge, 1999.
- [10] B. Sturmfels. *Gröbner Bases and Convex Polytopes*. American Mathematical Society University Lecture Series, **8**, Providence, RI, 1995.
- [11] B. Sturmfels and R. R. Thomas. Variation of cost functions in integer programming. *Mathematical Programming*, **77**(1997), pp. 357-387.
- [12] B. Sturmfels, N. V. Trung and W. Vogel. Bounds for degrees of projective schemes. *Mathematische Annalen*, **302**(1995), pp. 417-432.
- [13] R. R. Thomas. A Geometric Buchberger Algorithm for Integer Programming. *Mathematics of Operations Research*, **20**(1995), pp. 864-884.
- [14] R. Urbaniak, R. Weismantel and G. M. Ziegler. A variant of the Buchberger algorithm for integer programming. *SIAM Journal on Discrete Mathematics*, **10**(1997), pp. 96-108.
- [15] G. M. Ziegler. Gröbner bases and integer programming. In *Some Tapas of Computer Algebra* (A.M. Cohen, H. Cuypers and H. Sterk eds.), Springer-Verlag, Berlin, pp. 168-183, 1999.